

<b>Antibiotic Class</b>	<b>Contraindications</b>
Penicillins	History of hypersensitivity to penicillins, cephalosporins, imipenem (carbapenems)
Cephalosporins	Hypersensitivity to cephalosporins, imipenem (carbapenems), penicillins
Fluoroquinolones	Hypersensitivity to any of the fluoroquinolones; tendonitis or tendon rupture; use of QT interval-prolonging drugs ( <a href="#">amiodarone</a> , <a href="#">procainamide</a> )
Tetracyclines	Hypersensitivity to any of the tetracyclines
Macrolides	Hypersensitivity to any of the macrolides; treatment of meningitis (inadequate CNS penetration)
<a href="#">Vancomycin</a>	Hypersensitivity to <a href="#">vancomycin</a>
<a href="#">Clindamycin</a>	Hypersensitivity to <a href="#">clindamycin</a> ; treatment of meningitis (inadequate CNS penetration)
Aminoglycosides	Previous allergic or toxic reactions to any of the aminoglycosides
<a href="#">Metronidazole</a>	Hypersensitivity to <a href="#">metronidazole</a> ; first trimester of pregnancy
Sulfonamides	Hypersensitivity to sulfonamides or related drugs (sulfonylureas, thiazides, loop diuretics), pregnancy at term, lactation